

# A new semiempirical formula for exotic cluster decays of nuclei

M. Balasubramaniam, S. Kumarasamy and N. Arunachalam

Department of Physics, Manonmaniam Sundaranar University, Tirunelveli-627012, India.

Raj K. Gupta

Department of Physics, Panjab University, Chandigarh-160014, India.

(February 8, 2008)

A new semiempirical formula, with only three parameters, is proposed for cluster decay half-lives. The parameters of the formula are obtained by making a least squares fit to the available experimental data. The calculated half-lives are compared with an earlier proposed model-independent scaling law. Also, the calculated results of this formula are compared with the recent results of the preformed cluster model for  $^{12}\text{C}$  and  $^{14}\text{C}$  emissions from different deformed and superdeformed Nd and Gd parents. The results are in good agreement with experiments as well as other models.

23.90.+w, 25.85.Ca, 23.60.+e

## I. INTRODUCTION

In a radioactive decay series, the end product is reached not only via the emission of  $\alpha$  and  $\beta$  particles but also directly via heavy nuclei emission, the clusters, like Carbon, Oxygen, Fluorine, Neon, Magnesium and Silicon. Such a process is known as cluster decay, first proposed theoretically in 1980 by Săndulescu, Poenaru and Greiner [1], before its experimental realization in 1984 by Rose and Jones [2]. In this decay process a parent nucleus  $(A, Z)$ , with mass number  $A$  and charge number  $Z$ , breaks into two fragments, *viz.*, the emitted cluster  $(A_2, Z_2)$  and the associated daughter  $(A_1, Z_1)$ , where  $A=A_1+A_2$ ,  $Z=Z_1+Z_2$ . The light fragment  $(A_2, Z_2)$  is a cluster, heavier than the  $\alpha$ -particle but lighter than the lightest fission fragment observed so far.

The detection of cluster-decay was hindered mainly due to large pile-up of  $\alpha$  particles, but with improved facilities these difficulties are removed and a large number of cluster-decays are observed from different radioactive nuclei. In the years that followed, the cluster decay process has been studied extensively using different theoretical models with different realistic nuclear interaction potentials. In general, two kinds of models are used for explaining the observed and/or for predicting new decay modes. In one kind of these models, the  $\alpha$ -particle as well as the heavy cluster(s) were assumed to be pre-born in a parent nucleus, before they could penetrate the barrier with the available  $Q$ -value. These models are called the preformed cluster models [3–7]. In such a model, the clusters of different sizes (mass and/or charge numbers) are considered to be preformed in the parent nucleus, with

different probabilities. The Gamow-like barrier penetration is also taken into account in these models. In the other kind of models, only the Gamow's idea of barrier penetration is used, without considering the cluster(s) being or not being preformed in the parent nucleus. In other words, in this kind of models, called the unified fission models [1,8–13], the cluster radioactivity is considered as a simple barrier penetration phenomenon, in between the  $\alpha$ -decay and the spontaneous fission. In this paper, we attempt to give a model-independent, semiempirical formula for studying this above mentioned process of exotic cluster decay.

Geiger and Nuttall [14] were the first who proposed a semiempirical law connecting the  $\alpha$ -decay half-life and its  $Q$ -value. Now, a large number of  $\alpha$ -decays are observed from medium mass to superheavy nuclei, and several attempts have been made to give a universal formula [15–17]. These formulae vary among themselves mainly in the number of parameters. One such scaling law, proposed recently by Horoi *et al* [15], accounts for both the  $\alpha$  and cluster decays, and is given as,

$$\log T_{1/2} = (a_1 \mu^x + b_1) \left[ \frac{(Z_1 Z_2)^y}{\sqrt{Q}} - 7 \right] + (a_2 \mu^x + b_2). \quad (1)$$

This scaling law involves six parameters  $a_1 = 9.1$ ,  $b_1 = -10.2$ ,  $a_2 = 7.39$ ,  $b_2 = -23.2$ ,  $x = 0.416$  and  $y = 0.613$ , which were obtained by fitting 119  $\alpha$ -decays and 11 cluster-decays from various even-even parents. In the following, we propose a new semiempirical formula based on only three parameters that are least squares fitted to the cluster-decay data alone. So far, no attempt has been made to fit the  $\alpha$ -decay data.

The evolution of our proposed formula is presented in Section II, followed by a brief description of the preformed cluster model (PCM) of Gupta and collaborators [3–5] in Section III. The PCM is recently used by two of us and collaborators [18] to calculate the cluster-decay half-lives of some deformed and superdeformed Gd and Nd parents. Some of these results of Ref. [18] are used here for comparisons with the calculations based on the proposed new semiempirical formula. The results of our calculation are discussed in Section IV, and the summary and conclusions are presented in Section V.

## II. THE NEW SEMIEMPIRICAL FORMULA

We base the new formula for cluster decay half-lives on the following three simple experimental facts:

- (i) It is known from experiments that the cluster decay half-lives increase with the size (mass and/or charge) of the clusters. Hence, the empirical formula should contain terms showing direct dependence on the mass number and charge number of the cluster.
- (ii) The same cluster is emitted by different parents and hence the formula should contain dependence on the mass and charge asymmetries

$$\eta = \frac{A_1 - A_2}{A}; \quad \eta_z = \frac{Z_1 - Z_2}{Z}, \quad (2)$$

respectively.

- (iii) Since the  $\alpha$ -decay and cluster-decay are physically similar processes, the Q-dependence is taken to be the same as in Geiger-Nuttall law for  $\alpha$ -decay, i.e.,  $\log T_{1/2} \propto Q^{-1/2}$ .

Combining the above three results, we get

$$\log T_{1/2}^{AZ} = \frac{aA_2\eta + bZ_2\eta_z}{\sqrt{Q}} + c, \quad (3)$$

where the constants  $a = 10.603$ ,  $b = 78.027$  and  $c = -80.669$  are obtained by making a least squares fit of the available experimental half-lives for exotic cluster decays alone, with an rms deviation  $d_{rms}=0.89(s)$ , defined as  $d_{rms} = \sum_{i=1}^n \left[ \frac{y_i - f(x_i, a_j)}{\sigma_i} \right]^2$ , with  $f(x_i, a_j)$  denoting the function in Eq. (3),  $n$  the number of measurements and  $y_i$  the experimentally observed values. The  $\sigma_i^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$ , the variance, gives the standard deviation  $\sigma_i$ , with  $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$  giving the arithmetic average of the experimentally measured quantities. We refer to this formula as the *AZ-formula* (*AZF*). Note that the dependence on both  $\eta$  and  $\eta_z$  must be included in the formula (3) since these are separately measurable quantities, though the coupling between them is known to be weak [19–21], as is also evident when we consider the  $\eta$  and  $\eta_z$  dependence separately. These special cases of AZF formula, i.e.,  $b=0$  or  $a=0$  in (3), respectively, are expected to give reasonably good results, though poorer than the AZF results. These truncated expressions are referred to as the *AF* and *ZF* formulae in the following, whose constants are also obtained directly by the least squares fit to data. These are:  $a = 30.568$  and  $c = -51.348$  for AF and  $b = 112.197$  and  $c = -89.025$  for ZF, with rms deviations  $d_{rms}=1.652$  and  $1.112$  (s), respectively. Apparently, the rms deviations for truncated expressions are larger, and hence the fits are poorer, compared to the total expression (3). This will also be evident when we compare the results of these expressions with experimental data in Section IV.

## III. THE PREFORMED CLUSTER MODEL

In the preformed cluster model (PCM) of Gupta and collaborators [3–5], the decay half-life is given in terms of three factors, the  $P_0$ ,  $P$  and  $\nu$ , as

$$\log T_{1/2} = \ln 2 / P_0 P \nu, \quad (4)$$

where  $P_0$  is the preformation probability of the two fragments (the cluster and daughter nuclei) in their respective ground states,  $P$  the probability to tunnel the confining nuclear interaction barrier and  $\nu$  as an assault frequency.

For calculating  $P_0$  and  $P$ , Gupta *et al.* introduced the coupled motion in dynamical collective coordinates of mass asymmetry  $\eta$  and relative separation  $R$  via a stationary Schrödinger equation

$$H(\eta, R)\psi(\eta, R) = E\psi(\eta, R), \quad (5)$$

with the potential in it defined by the sum of experimental binding energies [22], the Coulomb and the proximity [23] potentials, as

$$V(\eta, R) = \sum_{i=1}^2 B_i(A_i, Z_i) + \frac{Z_1 Z_2 e^2}{R} + V_P. \quad (6)$$

Here the charges  $Z_i$  are fixed by minimizing the potential (without  $V_P$ ) in the charge asymmetry co-ordinate  $\eta_z$ . Equation (5) is solved in the decoupled approximation, which gives

$$P_0 \propto |\psi(\eta)|^2 \quad (7)$$

$$P \propto |\psi(R)|^2. \quad (8)$$

Only the ground state solution is relevant for the cluster decay to occur in the ground state of the daughter nucleus. Then, for  $\eta$ -motion, the properly normalized fractional preformation probability for, say, cluster  $A_2$  at a fixed  $R$  ( $=R_a=C_t=C_1+C_2$ ,  $C_i$  being the Süssmann central radii) is

$$P_0(A_2) = |\psi(\eta)|^2 \sqrt{B_{\eta\eta}} \frac{2}{A}. \quad (9)$$

For  $R$ -motion, the potential  $V(R)$  is obtained from Eq. (6) for fixed  $\eta$  and, instead of solving the corresponding radial Schrödinger equation, the WKB approximation is used for calculating the penetrability  $P$ . Finally, the assault frequency  $\nu$  in PCM is defined by considering that the total kinetic energy, shared between the two fragments, is the positive Q-value:

$$\nu = v/R_0 = \frac{\sqrt{2Q/mA_2}}{R_0}. \quad (10)$$

Here  $R_0$  is the spherical radius of the parent nucleus and  $m$ , the nucleon mass.

## IV. RESULTS AND DISCUSSION

Figure 1 and Table I give our calculated logarithms of decay half-lives by using the *AZ-formula* (3) for different clusters emitted from various radioactive parents, compared with the experimental data. In Fig. 1, we have also plotted the results of our calculations for *AF* ( $b = 0$ ) and *ZF* ( $a = 0$ ) versions of the *AZ-formula*. It is evident that the *AZF* fit to the data is better than the *AF* and *ZF* fits, as expected from our discussion above in Section II.

In Table I, we have also added the experimental data on *Q*-values and the results of another calculation using Eq. (1) of Horio *et al.* [15]. The comparison between the experiments and formulae (1) and (3) are also displayed in Fig. 2 for the illustrative cases of  $^{14}\text{C}$  and  $^{24}\text{Ne}$  cluster decays (respectively, the upper and lower panel). Apparently, our semiempirical *AZ-formula* is much closer to experiments, as compared to the other formula due to Horio *et al.* [15].

Finally, in order to compare the results of our semiempirical formula with the results of a model-dependent theoretical cluster decay calculation, we use a recent calculation [18] of two of us (MB and RKG) and collaborators based on PCM for the emission of  $^{12}\text{C}$  and  $^{14}\text{C}$  clusters from  $^{133-137}_{60}\text{Nd}$  and  $^{144-158}_{64}\text{Gd}$  parents. It may be mentioned here that these cluster-decay calculations are of interest only for nuclear structure information since their experimental observation may not be feasible in the near future [18]. Figs. 3(a) to (d) show the results of the PCM calculations, compared with those from the semiempirical *AZ-formula* (3) and the scaling law (1) of Horoi *et al.* [15]. We notice that, in general, the predictions of our semiempirical law lie higher than those of the PCM and Horoi *et al.* This is more so for Nd parents than for Gd parents where the predictions of the three calculations are nearly similar. The interesting point is that the three calculations predict an exactly the same structure for  $\log T_{1/2}$  vs. *A*, the parent mass number. For quantitative comparisons, it may be reminded here that the constants *a*, *b* and *c* of our semiempirical formula are obtained by fitting the data from trans-actinide region, which may or may not be good for this trans-tin region.

## V. SUMMARY

In this paper, we have proposed a model-independent three parameter formula for calculating the half-lives of cluster decays of nuclei. The evolution of the formula is based on three simple experimental observations about the characteristics of exotic cluster decays. The inputs of the formula are simply the mass and charge numbers of the parent and cluster, along with the *Q*-value of decay. The predictions of the proposed formula are comparable to another model-independent scaling law as well as to the well known preformed cluster model. The resulting

good comparisons suggest that this formula could be used to make predictions for the guide of new experiments, similar to that of Poenaru *et al.* [41].

## ACKNOWLEDGMENTS

One of the authors (M.B.) acknowledges with thanks the partial financial support by the Department of Science and Technology (DST), vide Grant NO. SR/FTP/PSA-02/2002. Also, the support by DST under the FIST programme vide letter No. SR/FST/PSI-005/2000 to the Department of Physics, M.S. University, Tirunelveli, India, is gratefully acknowledged.

## Figure Captions

Fig. 1 The logarithms of decay half-lives for different clusters emitted from various radioactive parents, calculated by using *AZ-formula* (*AZF*) and compared with experimental data. Also, the results of calculations for *AF* ( $b=0$ ) and *ZF* ( $a=0$ ) truncations of *AZF* are shown for comparisons. The parents are labelled at the top X-axis and the corresponding clusters emitted by these parents are labelled at the bottom X-axis.

Fig. 2 The experimental data on logarithms of decay half-lives for the emission of  $^{14}\text{C}$  and  $^{24}\text{Ne}$  clusters from different radioactive parents, compared with the results of calculations using *AZ-formula* (*AZF*) proposed here and the scaling law of Ref. [15].

Fig.3 The logarithms of half-lives for the emission of  $^{12,14}\text{C}$  clusters from different deformed and superdeformed Nd and Gd parents. Our calculations, using *AZ-formula* (*AZF*), are compared with the calculations based on PCM and the scaling law of Horoi *et al.* [15].

- 
- [1] A. Săndulescu, D.N. Poenaru and W. Greiner, *Sovt. J. Part. Nucl.* **11** (1980) 528.
- [2] H.J. Rose and G.A. Jones, *Nature* **307** (1984) 245.
- [3] R.K. Gupta, *Proceedings V<sup>th</sup> International Conference on Nuclear Reaction Mechanisms*, Varenna, Italy, 1988, ed. E. Gadioli (Ricerca Scientifica ed Educazione Permanente, Milano, 1988), p. 416.
- [4] S.S. Malik and R.K. Gupta, *Phys. Rev. C* **39** (1989) 1992.
- [5] S. Kumar and R.K. Gupta, *Phys. Rev. C* **55** (1997) 218.
- [6] R. Blendowske, T. Fliessbach and H. Walliser, *Nucl. Phys. A* **464** (1987) 75.
- [7] R. Blendowske and H. Walliser, *Phys. Rev. Lett.* **61** (1988) 1930.
- [8] Y.J. Shi and W.J. Swiatecki, *Phys. Rev. Lett.* **54** (1985) 300.
- [9] G.A. Pik-Pichak, *Sovt. J. Nucl. Phys.* **44** (1986) 923.
- [10] G. Shanmugam and B. Kamalaharan, *Phys. Rev. C* **38** (1988) 1377.
- [11] B. Buck and A.C. Merchant, *J. Phys. G: Nucl. Part. Phys.* **15** (1989) 615.
- [12] A. Săndulescu, R.K. Gupta, W. Greiner, F. Carstoiu, and M. Horoi, *Int. J. Mod. Phys. E* **1** (1992) 374.
- [13] G. Royer, R.K. Gupta and V.Yu. Denisov, *Nucl. Phys. A* **632** (1998) 275.
- [14] H. Geiger and J.M. Nuttall, *Phil. Mag.* **22** (1911) 613.
- [15] M. Horoi, B.A. Brown and A. Săndulescu, Preprint: <http://arxiv.org/archive/nucl-th/9403008>
- [16] A. Brown, *Phys. Rev. C* **46** (1992) 811.
- [17] G. Royer, *J. Phys. G: Nucl. Part. Phys.* **26** (2000) 1149.
- [18] R.K. Gupta, S. Dhaulta, R. Kumar, M. Balasubramanian, G. Münzenberg and W. Scheid, *Phys. Rev. C* **68** (2003) 034321.
- [19] J. Maruhn and W. Greiner, *Phys. Rev. Lett.* **32** (1974) 548.
- [20] R.K. Gupta, W. Scheid and W. Greiner, *Phys. Rev. Lett.* **35** (1975) 353.
- [21] R.K. Gupta, in *Heavy Elements and Related New Phenomena*, edited W. Greiner and R.K. Gupta, World Sc. Singapore, 1999, Vol. II, p.730.
- [22] G. Audi and A.H. Wapstra, *Nucl. Phys. A* **595** (1995) 4.
- [23] J. Blocki, J. Randrup, W.J. Swiatecki, and C.F. Tsang, *Ann. Phys. (N.Y)* **105** (1977) 427.
- [24] R. Bonetti, C. Chiesa, A. Guglielmetti, C. Migliorino, P. Monti, A.L. Pasinetti and H.L. Ravn, *Nucl. Phys. A* **576** (1994) 21.
- [25] P.B. Price, J.D. Stevenson, S.W. Barwick and H.L. Ravn, *Phys. Rev. Lett.* **54** (1985) 297.
- [26] E. Hourani, L. Rosier, G. Berrier-Ronsin, A. Elayi, A.C. Mueller, G. Rappenecker, G. Rotbard, G. Renou, A. Libe and L. Stab, *Phys. Rev. C* **44** (1991) 1424.
- [27] R. Bonetti, C. Chiesa, A. Guglielmetti, R. Matheoud, C. Migliorino, A.L. Pasinetti and H. L. Ravn, *Nucl. Phys. A* **562** (1993) 32.
- [28] E. Hourani, M. Hussonnois, L. Stab, L. Brillard, S. Gales and J.P. Schapira, *Phys. Lett. B* **160** (1985) 375.
- [29] R. Bonetti, C. Chiesa, A. Guglielmetti, C. Migliorino, A. Cesana and M. Terrani, *Nucl. Phys. A* **556** (1993) 115.
- [30] P. Qiangyan, Y. Weifan, Y. Shuanggui, L. Zongwei, M. Taotao, L. Yixiao, K. Dengming, Q. Jimin, L. Zihua, Z. Mutian and W. Shuhong, *Phys. Rev. C* **62** (2000) 044612.
- [31] P.B. Price, R. Bonetti, A. Guglielmetti, C. Chiesa, R. Matheoud, C. Migliorino and K.J. Moody, *Phys. Rev. C* **46** (1992) 1939.
- [32] S.P. Tretyakova, A. Săndulescu, V.L. Mikheev, D. Hasegan, I.A. Lebedev, Yu.S. Zamyatnin, Yu.S. Korotkin and B. F. Myasoedov, *JINR Dubna Rapid Commun.* **13** (1985) 34.
- [33] A. Săndulescu, Yu.S. Zamyatnin, I.A. Lebedev, B. F. Myasoedov, S.P. Tretyakova and D. Hasegan, *JINR Dubna Rapid Commun.* **15** (1984) 5.
- [34] S.W. Barwick, P.B. Price and J.D. Stevenson, *Phys. Rev. C* **31** (1985) 1984.
- [35] R. Bonetti, C. Chiesa, A. Guglielmetti, C. Migliorino, A. Cesan, M. Terrani and P. B. Price, *Phys. Rev. C* **44** (1991) 888.
- [36] K.J. Moody, E.K. Hulet, S. Wang and P. B. Price, *Phys. Rev. C* **39** (1989) 2445.
- [37] S.P. Tretyakova, Yu.S. Zamyatnin, V.N. Kovantsev, Yu. S. Korotkin, V. L. Mikheev and G. A. Timofeev, *Z. Phys. A* **33** (1989) 349.
- [38] A.A. Oglobin, N.Z. Venikov, S.K. Lisin, S.V. Pirozhkov, V.A. Pchelin, Yu.F. Rodionov, V.M. Semochkin, V.A. Shabrov, I.K. Shvetsov, V.M. Shubko, S.P. Tretyakova and V.L. Mikheev, *Phys. Lett. B* **235** (1990) 35.
- [39] S.P. Tretyakova, V.L. Mikheev, V.A. Ponomarenko, A.N. Golovchenko, A.A. Oglobin and V.A. Shigin, *JETP Lett.* **59** (1994) 394.
- [40] S. Wang, D. Snowden-Ifft, P.B. Price, K. J. Moody and E. K. Hulet, *Phys. Rev. C* **39** (1989) 1647.
- [41] D.N. Poenaru, D.Schnabel, W. Greiner, D. Mazilu and R. Gherghescu, *At. Data Nucl. Data Tables* **48** (1991) 231.

TABLE I. The logarithms of decay half-lives of different clusters emitted from various radioactive nuclei, calculated by using the semiempirical *AZ-formula* (3) and the scaling law (1) compared with the experimental data. The experimental Q-values are also given and the data are tabulated in the increasing order of the mass number of the clusters. The last column gives the reference to the source of experimental data.

| Parent            | Cluster          | Daughter          | $Q^{Expt.}$ (MeV) | $\log T_{1/2}^{AZF}$ (s) | $\log T_{1/2}^{Ref.[15]}$ (s) | $\log T_{1/2}^{Expt.}$ (s) | Ref.    |
|-------------------|------------------|-------------------|-------------------|--------------------------|-------------------------------|----------------------------|---------|
| <sup>221</sup> Fr | <sup>14</sup> C  | <sup>207</sup> Tl | 31.28             | 14.54                    | 13.56                         | 14.52                      | [24]    |
| <sup>221</sup> Ra | <sup>14</sup> C  | <sup>207</sup> Pb | 32.39             | 12.96                    | 12.28                         | 13.39                      | [24]    |
| <sup>222</sup> Ra | <sup>14</sup> C  | <sup>208</sup> Pb | 33.05             | 11.98                    | 11.00                         | 11.01                      | [25]    |
| <sup>223</sup> Ra | <sup>14</sup> C  | <sup>209</sup> Pb | 31.85             | 13.81                    | 13.38                         | 15.20                      | [25]    |
| <sup>224</sup> Ra | <sup>14</sup> C  | <sup>210</sup> Pb | 30.54             | 15.94                    | 16.13                         | 15.68                      | [26]    |
| <sup>225</sup> Ac | <sup>14</sup> C  | <sup>211</sup> Bi | 30.48             | 16.20                    | 17.26                         | 17.16                      | [27]    |
| <sup>226</sup> Ra | <sup>14</sup> C  | <sup>212</sup> Pb | 28.21             | 20.08                    | 21.50                         | 21.19                      | [28]    |
| <sup>228</sup> Th | <sup>20</sup> O  | <sup>208</sup> Pb | 44.72             | 21.90                    | 21.20                         | 20.72                      | [29]    |
| <sup>230</sup> U  | <sup>22</sup> Ne | <sup>208</sup> Pb | 61.59             | 21.78                    | 19.28                         | 20.14                      | [30]    |
| <sup>231</sup> Pa | <sup>23</sup> F  | <sup>208</sup> Pb | 51.84             | 24.30                    | 23.85                         | 26.02                      | [31]    |
| <sup>230</sup> Th | <sup>24</sup> Ne | <sup>206</sup> Hg | 57.78             | 25.77                    | 23.88                         | 24.61                      | [32]    |
| <sup>231</sup> Pa | <sup>24</sup> Ne | <sup>207</sup> Tl | 60.42             | 23.62                    | 21.30                         | 23.23                      | [33]    |
| <sup>232</sup> U  | <sup>24</sup> Ne | <sup>208</sup> Pb | 62.31             | 22.24                    | 19.94                         | 21.08                      | [34]    |
| <sup>233</sup> U  | <sup>24</sup> Ne | <sup>209</sup> Pb | 60.5              | 23.87                    | 22.53                         | 24.83                      | [32]    |
| <sup>234</sup> U  | <sup>24</sup> Ne | <sup>210</sup> Pb | 58.84             | 25.42                    | 25.01                         | 25.92                      | [35,36] |
| <sup>235</sup> U  | <sup>24</sup> Ne | <sup>211</sup> Pb | 57.36             | 26.87                    | 27.31                         | 27.42                      | [35,37] |
| <sup>233</sup> U  | <sup>25</sup> Ne | <sup>208</sup> Pb | 60.75             | 24.15                    | 22.98                         | 24.83                      | [32]    |
| <sup>235</sup> U  | <sup>25</sup> Ne | <sup>210</sup> Pb | 57.83             | 26.94                    | 27.49                         | 27.42                      | [35,37] |
| <sup>234</sup> U  | <sup>26</sup> Ne | <sup>208</sup> Pb | 59.47             | 25.85                    | 25.75                         | 25.92                      | [35,36] |
| <sup>234</sup> U  | <sup>28</sup> Mg | <sup>206</sup> Hg | 74.13             | 26.24                    | 24.74                         | 27.54                      | [37]    |
| <sup>236</sup> Pu | <sup>28</sup> Mg | <sup>208</sup> Pb | 79.67             | 23.01                    | 20.83                         | 21.67                      | [38]    |
| <sup>236</sup> U  | <sup>28</sup> Mg | <sup>208</sup> Hg | 71.69             | 28.18                    | 27.98                         | 27.58                      | [39]    |
| <sup>238</sup> Pu | <sup>28</sup> Mg | <sup>210</sup> Pb | 75.93             | 25.70                    | 25.39                         | 25.70                      | [40]    |
| <sup>236</sup> U  | <sup>30</sup> Mg | <sup>206</sup> Hg | 72.51             | 28.36                    | 28.36                         | 27.58                      | [39]    |
| <sup>238</sup> Pu | <sup>30</sup> Mg | <sup>208</sup> Pb | 77.03             | 25.71                    | 25.41                         | 25.70                      | [40]    |
| <sup>238</sup> Pu | <sup>32</sup> Si | <sup>206</sup> Hg | 91.21             | 25.99                    | 25.68                         | 25.27                      | [40]    |




Fig.1 "A new semiempirical formula for..." by M. Balasubramaniam et al.

Fig.2 "A new semiempirical formula for..." by M. Balasubramaniam et al.




Fig.3 "A new semiempirical formula for..." by M. Balasubramaniam et al.

